

Seasonal Flu Criteria with further detail:

Or any patient (aged 6 months and over) with:

Chronic respiratory disease	including asthma requiring continuous or repeated use of inhaled or systemic steroids or with previous exacerbations requiring hospitalisation) and children who have previously been admitted to hospital for lower respiratory tract disease.
Chronic heart disease	Congenital heart disease, hypertension with cardiac complications, chronic heart failure, individuals requiring regular medication and/or follow-up for ischaemic heart disease.
Chronic renal disease	Chronic kidney disease at stage 3, 4 or 5, chronic kidney failure, nephrotic syndrome, kidney transplantation.
Chronic liver disease	Cirrhosis, biliary artesia, chronic hepatitis
Chronic neurological disease	Stroke, TIA. Clinicians should consider on an individual basis the clinical needs of patients including individuals with cerebral palsy, multiple sclerosis and related or similar conditions; or hereditary and degenerative disease of the nervous system or muscles; or severe neurological disability.
Diabetes	Type 1 diabetes, type 2 diabetes requiring insulin or oral hypoglycaemic drugs, diet-controlled diabetes.
Immunosuppression	Immunosuppression due to disease or treatment. Patients undergoing chemotherapy leading to immunosuppression. Asplenia or splenic dysfunction, HIV infection at all stages.
Pregnant ladies	Pregnant women at any stage of pregnancy (first, second or third trimesters).